

# Certified Urologic Associate Practice Test Questions and Answers

**1. What is the normal range for a post-void residual (PVR) measurement in adults?**

- A) 150-200 mL
- B) Less than 50 mL
- C) 100-150 mL
- D) 200-300 mL

**2. Which diagnostic procedure is considered the gold standard for evaluating bladder function and urodynamics?**

- A) Simple cystoscopy
- B) Multichannel urodynamic study
- C) Ultrasound of the bladder
- D) Urine dipstick test

**3. What is the most common cause of acute urinary retention in older men?**

- A) Urinary tract infection
- B) Benign prostatic hyperplasia (BPH)
- C) Bladder cancer
- D) Kidney stones

**4. During cystoscopy preparation, which patient instruction is most critical for procedure safety?**

- A) Drink extra fluids the night before
- B) Report any signs of urinary tract infection or fever before the procedure
- C) Take aspirin for pain prevention
- D) Avoid all medications for 48 hours prior

Answers: 1-B 2-B 3-B 4-B

For More Certified Urologic Associate Questions and Answers FREE, Certified Urologic Associate Online Prep Training, Certified Urologic Associate Exam, Certified Urologic Associate Study Guide, Certified Urologic Associate Flashcards, Certified Urologic Associate Quizzes visit:

**Certified Urologic Associate Practice Test**

Practice Test Geeks © All Rights Reserved